

Surgical Site Infection Rates in Elective vs Emergency Laparotomy

Campuzan Maya¹, Guillermo Latorro², Johnayro Gutirre³

Abstract Introduction: Surgical site infections (SSIs) remain one of the most common postoperative complications following abdominal surgeries. Laparotomy, particularly when performed under emergency conditions, is associated with increased morbidity due to contamination, delayed presentation, and physiological instability. **Materials and Methods:** A prospective comparative observational study was conducted over 18 months in a tertiary care hospital. A total of 200 patients undergoing laparotomy were enrolled and divided into two groups: Elective (n=100) and Emergency (n=100). Demographic variables, operative details, wound classification, and postoperative outcomes were recorded. SSIs were diagnosed based on CDC criteria. Statistical analysis was performed using chi-square test and logistic regression.

Results: Overall SSI rate was 18%. Elective group showed 8% SSI rate, whereas Emergency group demonstrated 28% SSI rate ($p < 0.001$). Risk factors significantly associated with SSI included emergency surgery, contaminated wounds, operative duration > 2 hours, anemia, diabetes mellitus, and delayed antibiotic administration. **Conclusion:** Emergency laparotomy is associated with significantly higher SSI rates compared to elective laparotomy. Optimization of patient factors, strict aseptic technique, timely antibiotics, and perioperative monitoring can reduce infection rates.

Keywords: Surgical site infection, Laparotomy, Emergency surgery, Elective surgery, Postoperative complications, Abdominal surgery.

¹ Clinical Hematology Laboratory, University of Antioquio.

² Clinical Hematology Laboratory, University of Antioquio.

³ Clinical Hematology Laboratory, University of Antioquio.



INTRODUCTION

Surgical site infection (SSI) is defined as infection occurring within 30 days after a surgical procedure involving skin, subcutaneous tissue, deep soft tissue, or organ space¹. SSIs account for nearly 20% of healthcare-associated infections globally and significantly contribute to prolonged hospital stay, increased healthcare costs, and patient morbidity². Despite advancements in antiseptic protocols, antimicrobial prophylaxis, and surgical techniques, SSI remains a persistent challenge in abdominal surgeries³.

Laparotomy, a surgical incision into the abdominal cavity, is commonly performed for various conditions such as gastrointestinal obstruction, malignancies, perforation peritonitis, trauma, and inflammatory diseases⁴. The risk of SSI in laparotomy is influenced by several factors including wound contamination level, duration of surgery, patient comorbidities, nutritional status, and emergency nature of the procedure⁵.

Elective laparotomies are performed under controlled conditions after adequate preoperative optimization of patients. Proper bowel preparation, prophylactic antibiotics, glycemic control, and sterile surgical environments contribute to lower infection rates⁶. In contrast, emergency laparotomies are often conducted in patients with hemodynamic instability, sepsis, bowel perforation, or trauma, where contamination risk is significantly higher⁷.

The World Health Organization reports that SSI incidence ranges from 1-10% in developed countries but may reach 30% in low- and middle-income countries⁸. Studies have consistently demonstrated that emergency surgeries carry a 2-4 times higher risk of SSI compared to elective

procedures⁹. The presence of fecal contamination, delayed presentation, and inadequate physiological preparation are major contributing factors¹⁰.

Host-related factors such as diabetes mellitus, anemia, obesity, hypoalbuminemia, smoking, and immunosuppression further increase susceptibility to postoperative infections¹¹. Additionally, prolonged operative time (>2 hours) and higher American Society of Anesthesiologists (ASA) scores correlate strongly with SSI incidence¹².

Understanding the differential risk between elective and emergency laparotomy is essential for implementing targeted preventive strategies. Enhanced Recovery After Surgery (ERAS) protocols, standardized antibiotic timing, wound protectors, and negative pressure wound therapy have shown promising reductions in SSI rates¹³.

Given the burden of SSI and its impact on surgical outcomes, this study aims to compare the incidence and risk factors of SSI in elective versus emergency laparotomy patients in a tertiary care setting.

MATERIALS AND METHODS

Prospective comparative observational study conducted over 18 months in a tertiary care hospital.

Sample Size

200 patients undergoing laparotomy.
Elective group: 100 patients
Emergency group: 100 patients

Inclusion Criteria

- Age ≥ 18 years
- Patients undergoing midline laparotomy
- Both genders
- Informed consent obtained

- HIV-positive patients

Exclusion Criteria

- Laparoscopic surgeries
- Re-laparotomy within 30 days
- Patients on immunosuppressive therapy
- Patients with pre-existing wound
- Pediatric patients

Data Collection

Demographic data: Age, gender, BMI

Clinical parameters: Comorbidities (DM, HTN), hemoglobin, serum albumin

Operative variables: Duration of surgery, wound classification (clean, clean-contaminated, contaminated, dirty)

Antibiotic timing: Within 60 minutes before incisión infection

Parameters Studied

Based on CDC criteria:

- Superficial incisional
 - Deep incisional
 - Organ-space infection
- Follow-up: 30 days postoperatively.

Statistical Analysis

- Chi-square test
- Logistic regression
- p-value <0.05 considered significant

Ethical approval obtained from Institutional Ethics Committee.

RESULTS

Table 1: Demographic Profile

| Variable | Elective (n=100) | Emergency (n=100) |
|--------------|------------------|-------------------|
| Mean Age | 45.2 ±12 | 48.6 ±14 |
| Male (%) | 58 | 62 |
| Diabetes (%) | 18 | 26 |
| Anemia (%) | 22 | 40 |

Emergency patients had higher prevalence of anemia and diabetes.

Table 2: Wound Classification

| Classification | Elective | Emergency |
|--------------------|----------|-----------|
| Clean | 40 | 5 |
| Clean-contaminated | 45 | 20 |
| Contaminated | 10 | 40 |
| Dirty | 5 | 35 |

Majority of emergency surgeries were contaminated/dirty wounds.

Table 3: Duration of Surgery

| Duration | Elective | Emergency |
|----------|----------|-----------|
| <2 hours | 70 | 42 |
| >2 hours | 30 | 58 |

Emergency procedures had significantly longer duration.

Table 4: SSI Incidence

| Group | SSI Present | SSI Absent | Rate |
|-----------|-------------|------------|------|
| Elective | 8 | 92 | 8% |
| Emergency | 28 | 72 | 28% |
| p <0.001 | | | |

Emergency laparotomy showed significantly higher SSI rate.

Table 5: Type of SSI

| Type | Elective | Emergency |
|-------------|----------|-----------|
| Superficial | 6 | 15 |
| Deep | 2 | 7 |
| Organ-space | 0 | 6 |

Organ-space infections were more common in emergency cases.

Table 6: Risk Factor Analysis

| Risk Factor | Odds Ratio | p-value |
|--------------------|------------|---------|
| Emergency surgery | 3.8 | <0.001 |
| Contaminated wound | 4.2 | <0.001 |
| Duration >2h | 2.6 | 0.01 |
| Diabetes | 2.1 | 0.03 |
| Anemia | 2.8 | 0.01 |

Emergency surgery and contaminated wounds were strongest predictors.

DISCUSSION

The present study demonstrates that SSI rates are significantly higher in emergency laparotomy (28%) compared to elective laparotomy (8%). These findings align with global data reporting SSI rates ranging from 20-35% in emergency abdominal surgeries¹⁴.

The higher incidence in emergency cases is attributable to increased contamination levels, hemodynamic instability, and inadequate preoperative optimization¹⁵. Contaminated and dirty wounds showed a fourfold increase in infection risk, similar to findings by Smith et al.¹⁶.

Operative duration exceeding two hours was another independent predictor. Prolonged exposure increases bacterial load and tissue trauma¹⁷. Diabetes mellitus doubled the risk of SSI, consistent with findings from WHO multicenter studies¹⁸.

Organ-space infections were predominantly seen in emergency surgeries, likely due to perforation peritonitis and intra-abdominal sepsis¹⁹. Early antibiotic administration and improved perioperative glycemic control have shown effectiveness in reducing SSI rates²⁰.

Enhanced recovery protocols, wound protectors, and negative pressure

dressings have demonstrated significant reductions in high-risk abdominal surgeries²¹. Resource-limited settings require implementation of standardized infection prevention bundles²².

Overall, this study reinforces the need for risk stratification and aggressive preventive measures in emergency laparotomy patients.

CONCLUSION

Emergency laparotomy carries significantly higher risk of surgical site infection compared to elective procedures. Wound contamination, prolonged surgery, anemia, and diabetes are major contributing factors. Implementation of evidence-based preventive strategies and optimization of patient comorbidities are essential to reduce SSI burden.

REFERENCES

1. Allegranzi B, Bischoff P, de Jonge S, Kubilay NZ, Zayed B, Gomes SM, et al. New WHO recommendations on preoperative measures for surgical site infection prevention: an evidence-based global perspective. *Lancet Infect Dis*. 2016;16(12):e276–87.
2. Berríos-Torres SI, Umscheid CA,

- Bratzler DW, Allen B, Stone EC, Kelz RR, et al. Centers for Disease Control and Prevention guideline for the prevention of surgical site infection, 2017. *JAMA Surg.* 2017;152(8):784–91.
3. Ban KA, Minei JP, Laronga C, Harbrecht BG, Jensen EH, Fry DE, et al. American College of Surgeons and Surgical Infection Society guidelines for the prevention of surgical site infection. *Surg Infect (Larchmt).* 2017;18(4):379–82.
 4. GlobalSurg Collaborative. Surgical site infection after gastrointestinal surgery in high-income, middle-income, and low-income countries: a prospective, international cohort study. *Lancet Infect Dis.* 2018;18(5):516–25.
 5. World Health Organization. Global guidelines for the prevention of surgical site infection, 2nd edition. Geneva: WHO; 2018.
 6. Badia JM, Casey AL, Petrosillo N, Hudson PM, Mitchell SA, Crosby C. Impact of surgical site infection on healthcare costs and patient outcomes: a systematic review. *J Hosp Infect.* 2017;96(1):1–15.
 7. Martin ET, Kaye KS, Knott C, Nguyen H, Santarossa M, Evans R, et al. Diabetes and risk of surgical site infection: a systematic review and meta-analysis. *Clin Infect Dis.* 2016;59(5):543–50.
 8. Cheng H, Chen BP, Soleas IM, Ferko NC, Cameron CG, Hinoul P. Prolonged operative duration and surgical site infection risk: a meta-analysis. *Ann Surg.* 2017;265(1):6–16.
 9. Anderson DJ, Podgorny K, Berríos-Torres SI, Bratzler DW, Dellinger EP, Greene L, et al. Strategies to prevent surgical site infections in acute care hospitals. *Infect Control Hosp Epidemiol.* 2019;40(4):409–32.
 10. De Simone B, Sartelli M, Coccolini F, Ball CG, Brambillasca P, Chiarugi M, et al. Intra-abdominal infections: WSES guidelines update. *World J Emerg Surg.* 2020;15:38.
 11. Bhangu A, Nepogodiev D, Glasbey JC, Li E, Omar OM, Soreide K, et al. Mortality of emergency abdominal surgery in low-income and middle-income countries. *Lancet.* 2018;391(10130):139–48.
 12. Tanner J, Dumville JC, Norman G, Fortnam M. Surgical hand antisepsis to reduce surgical site infection. *Cochrane Database Syst Rev.* 2016;1:CD004288.
 13. Hyldig N, Vinter CA, Kruse M, Mogensen O, Bille C, Sorensen JA, et al. Prophylactic incisional negative pressure wound therapy reduces surgical site infection. *Ann Surg.* 2019;269(4):664–70.
 14. Pinkney TD, Bartlett DC, Hawkins W, Mak T, Nepogodiev D, Sarr MG, et al. Reduction of surgical site infection using wound-edge protection devices: randomized trial. *Lancet.* 2019;393(10171):1263–70.
 15. Kougiass P, Sharath SE, Barshes NR. Risk factors for surgical site infection in emergency general surgery. *Surgery.* 2017;162(4):715–23.
 16. Ingraham AM, Cohen ME, Bilimoria KY, Ko CY, Hall BL. Comparison of outcomes after emergency versus elective colorectal surgery. *Ann Surg.* 2018;268(4):603–9.
 17. De Lissovoy G, Fraeman K, Hutchins V, Murphy D, Song D, Vaughn BB. Surgical site infection: incidence and impact on hospital utilization and treatment costs. *Am J Infect Control.* 2016;44(12):1463–9.
 18. Ljungqvist O, Scott M, Fearon KC. Enhanced recovery after surgery: a review. *JAMA Surg.* 2017;152(3):292–8.
 19. GlobalSurg Collaborative. Determinants of morbidity and mortality following emergency abdominal surgery. *Br J Surg.* 2019;106(8):1046–56.



20. Allegranzi B, Zayed B, Bischoff P, Kubilay NZ, de Jonge S, de Vries F, et al. New WHO recommendations on intraoperative and postoperative measures for SSI prevention. *Lancet Infect Dis.* 2016;16(12):e288–303.
21. Young PY, Khadaroo RG. Surgical site infections. *Surg Clin North Am.* 2019;99(5):927–41.
22. Sartelli M, Chichom-Mefire A, Labricciosa FM, Hardcastle T, Abu-Zidan FM, Adesunkanmi AK, et al. The management of intra-abdominal infections worldwide. *World J Emerg Surg.* 2017;12:29.
23. Cheng K, Robertson C, Murray A, Norman G. Wound irrigation and SSI prevention. *J Hosp Infect.* 2017;95(2):135–43.
24. Teshome BF, Desalegn B, Fentie M. Surgical site infection and associated factors among emergency laparotomy patients. *BMC Surg.* 2020;20:156.
25. Watanabe M, Suzuki H, Nomura S, Yamada K, Kobayashi M. Risk factors for surgical site infection in abdominal surgery: a multicenter study. *Int J Surg.* 2018;54(Pt A):165–9.